

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89118014.3

(51) Int. Cl.⁵: **C07D 215/42 , A61K 31/47 ,
C07D 213/74 , A61K 31/44**

(22) Date of filing: 28.09.89

(30) Priority: 30.09.88 JP 246237/88

(43) Date of publication of application:
04.04.90 Bulletin 90/14

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

(71) Applicant: **Chugai Seryaku Kabushiki Kaisha**
5-1, 5-chome, Ukima Kita-ku
Tokyo(JP)


(72) Inventor: **Miura, Yutaka**
Nochi Mansions B-201 10-20, Kobara-cho
1-chome
Mishima-shi Shizuoka-ken(JP)
Inventor: **Yoshida, Mitsutaka**

24-2, Fujimidai
Mishima-shi Shizuoka-ken(JP)
Inventor: **Fujimura, Yasuo**
2-94-8, Sakae, Ina-cho
Kitaadachi-gun Saitama-ken(JP)
Inventor: **Takaku, Sakae**
381-12, Oazakoizumi
Ageo-shi Saitama-ken(JP)
Inventor: **Noda, Yukifumi**
888-24, Shimotogari Nagaizumi-cho
Sunto-gun Shizuoka-ken(JP)

(74) Representative: **Vossius & Partner**
Siebertstrasse 4 P.O. Box 86 07 67
D-8000 München 86(DE)

(54) **Novel 3,4-diaminoquinoline and pyridine compounds.**


(57) 4-aminopyridine derivatives represented by the general formula:



[where R₁ is a hydrogen atom, a hydroxyl group, a linear or branched lower alkyl group or a cycloalkyl group which may be substituted by a hydroxyl group, a lower alkoxy group, a lower alkyl or a cycloalkyl group which contains a carbonyl group, a morpholino group or a group -NH-B (where B is a lower alkyl group, a cycloalkyl group or a phenyl group); R₂ and R₃ which may be the same or different each represents a hydrogen atom, a lower alkyl group or a loweralkylcarbonyl group, or when taken together, form an azacycloalkyl group, a morpholino group or an N-methylpiperazinyl group together with the nitrogen atom; R₄ and R₅ each represents a hydrogen atom, or when taken together with the ring A, form a quinoline ring or a 5,6,7,8-tetrahydroquinoline ring, provided that when each of R₁, R₄ and R₅ is a hydrogen atom, R₂ and R₃ are neither a hydrogen atom nor a methyl group at the same time, and when R₁ is a hydrogen atom and R₄ and R₅ taken together with the ring A form a quinoline ring, R₂ and R₃ are neither a hydrogen atom nor an ethyl group at the same time], and an acid addition salts of said 4-aminopyridine derivatives. These compounds are useful as active ingredients in pharmaceutical compositions for improving psychoneural function, especially in the treatment of Alzheimer type dementia and promotion of mnemonic and learning performance.

NOVEL 3,4-DIAMINOQUINOLINE AND PYRIDINE COMPOUNDS

The present invention relates to 4-aminopyridine derivatives represented by the general formula (I):



where R_1 is a hydrogen atom, a hydroxyl group, a linear or branched lower alkyl group or a cycloalkyl group which may be substituted by a hydroxyl group, a lower alkoxy group, a lower alkyl or a cycloalkyl group which contains a carbonyl group, a morpholino group or a group $-NH-B$ (where B is a lower alkyl group, a cycloalkyl group or a phenyl group); R_2 and R_3 which may be the same or different each represents a hydrogen atom, a lower alkyl group or a loweralkylcarbonyl group, or when taken together, form an azacycloalkyl group, a morpholino group or an N-methylpiperazinyl group together with the nitrogen atom; R_4 and R_5 each represents a hydrogen atom, or when taken together with the ring A , form a quinoline ring or a 5,6,7,8-tetrahydroquinoline ring, provided that when each of R_1 , R_4 and R_5 is a hydrogen atom, R_2 and R_3 are neither a hydrogen atom nor a methyl group at the same time, and when R_1 is a hydrogen atom and R_4 and R_5 taken together with the ring A form a quinoline ring, R_2 and R_3 are neither a hydrogen atom nor an ethyl group at the same time. The present invention also relates to acid addition salts of said 4-aminopyridine derivatives.

With the progress of "aging society", the population of aged persons has recently increased and the increase in the number of senile diseases is correspondingly rapid. Among various senile diseases, dementia is of particular importance since the mechanism of its occurrence has not been fully unravelled and it can be fatal depending on its severity. Under these circumstances, the advent of an effective therapeutic for senile dementia is strongly desired.

Numerous studies on senile dementia have been conducted to date. The phenomena that have been reported to occur in patients with senile dementia include impairment of the central cholinergic nervous function on account of decreases in the number of choline acetyltransferase (CAT) and acetylcholinesterase (AChE) which are enzymes that synthesize and decompose, respectively, acetylcholine known to be the transmitter of cholinergic neurons (see British Medical Journal, 2: 1457 - 1459, 1978; Brain, 107: 507 - 518, 1984; Journal of the Neurological Sciences, 57: 407 - 417, 1982; and Lancet, 2: 1403, 1976) and impairment of the central noradrenergic nervous function (see British Medical Journal, 282: 93-94, 1981). Researchers are making active efforts to develop drugs that are suitable for the symptomatic treatment of these phenomena.


It is the object of the present invention to provide compounds which are therapeutically effective in reactivating the nervous function impaired by senile dementia. This object was achieved by the surprising finding that the compounds represented by the general formula (I) exhibited unique pharmacological effects in that they activated the nervous function of animals and accelerated their mnemonic and learning performance.

Fig. 1 shows mean correct response vs the dose of a compound of the present invention administered to rats under a radial-arm maze task in Test Example 6.

The terms "lower alkyl" and "lower alkoxy", taken alone or as part of the larger group, designate residues having 1 to 8, preferably 1 to 5 carbon atoms. The term "cycloalkyl" designates residues having 3 to 8, preferably 3 to 6 carbon atoms.

The compounds of the general formula (I) are novel and may be prepared by the following methods (a) - (h), the choice of which depends on R_1 in the general formula (I).


(a) If R_1 is a branched lower alkyl group or a cycloalkyl group which are substituted by a hydroxyl group, or if it is a lower alkyl group or a cycloalkyl group which contain a carbonyl group, the following procedure is taken: a starting compound represented by the general formula (II):



(II)

(where R_A and R_B each represents a hydrogen atom or, when taken together with the ring A, form a quinoline ring; X is a halogen) is reacted with an enamine corresponding to the end compound, thereby introducing a substituent on both 2 and 3 positions, and subsequently, the nitro group on 4 position is reduced, optionally followed by further reduction. The reaction with enamine is performed within an inert solvent such as chloroform or methylene chloride, and the enamine to be used corresponds to the end compound as exemplified by 1-(N-morpholino)cyclohexene. The reaction temperature ranges from 20 to 80°C and is preferably room temperature. The reduction of the nitro group on 4 position involves an ordinary reduction reaction and may be accomplished by hydrogenation in the presence of a catalyst such as palladium-on-carbon, Raney nickel or Raney cobalt. The reduction reaction is performed in an inert solvent such as tetrahydrofuran or an alcohol. A compound of the general formula (1) wherein R_4 and R_5 combine with the pyridine ring to form a 5,6,7,8-tetrahydroquinoline ring is obtained by performing further reduction in an acidic solvent such as trifluoroacetic acid or acetic acid in the presence of a reduction catalyst such as platinum dioxide or Raney nickel.


(b) If R_1 is a branched lower alkyl group or a cycloalkyl group, a compound of the general formula (II) is reacted with an enamine and the nitro group on 4 position is reduced as in (a). The resulting compound of the general formula (III):



(III)


(where R_C is a lower alkyl group or a cycloalkyl group which contain a carbonyl group; R_A , R_B , R_2 and R_3 are each the same as defined above) is reacted with a compound of the general formula $Y-NHNH_2$ (where Y is a paratoluenesulfonyl group or a hydrogen atom) in a solvent such as alcohol at 20 - 80°C, preferably at room temperature. By subsequent reaction with a reducing agent such as $NaBH_4$, $LiAlH_4$ or BH_3 , a compound of the general formula (I) is obtained. By performing further reduction which is optional, a 5,6,7,8-tetrahydroquinoline based end compound can be obtained.

(c) If R_1 is $-\text{NH}-\text{C}_6\text{H}_5$ or $-\text{NH}-\text{C}_6\text{H}_{11}$, the following scheme may be employed:




(where R_2 , R_3 , R_A , R_B and X have the same meanings as defined above: C_6H_5 denotes a phenyl or cyclohexyl group).

The conversion from the general formula (II) to (IV) is carried out by reacting the compound (II) with a corresponding amine in an inert solvent such as tetrahydrofuran or an alcohol, preferably at room temperature. The conversion from the general formula (IV) to (V) is carried out using phenyl isocyanate in a solvent such as dimethylformamide or dimethyl acetamide. The reaction temperature is in the range of 50 - 100°C, preferably 80°C. The conversion from the general formula (V) to (VI) involves an ordinary step of reduction and may be accomplished by hydrogenation in the presence of a catalyst such as palladium-on-carbon, Raney nickel or Raney cobalt. The reduction reaction is performed in an inert solvent such as tetrahydrofuran or an alcohol, preferably at room temperature. By further reducing the compound of the general formula (VI) with a suitable catalyst such as platinum dioxide or Raney cobalt, an end compound wherein R_1 is



or a 5,6,7,8-tetrahydroquinoline based compound is obtained (general formula (VII)).

(d) If R_1 is $-\text{NH}-\text{RD}$ (RD is a lower alkyl group), the following scheme may be employed:




(where R_2 , R_3 , R_A , R_B , R_D and X are the same as defined above).

The conversion from the general formula (II) to (IV) is carried out as in (c). The conversion from the

general formula (IV) to (VIII) is carried out by reacting the compound of formula (IV) with RD-NCO (where RD is a lower alkyl group) in a sealed tube. The reaction is performed in an inert solvent such as dimethylformamide at 100 - 150 °C, preferably at 130 °C, for a period of 4 - 10 hours, preferably 6 hours. Conversion from the general formula (VIII) to (IX) may be accomplished by reducing the nitro group as in


(a) - (c). If desired, further reduction may be performed by treatment with a suitable catalyst such as platinum dioxide or Raney cobalt and this leads to the production of an end compound of the general formula (X).

(e) If R₁ is a lower alkoxy group, the following scheme may be employed:



The conversion from the general formula (II) to (IV) is carried out as in (c). The conversion from the general formula (IV) to (XI) may be carried out in the usual manner and an α -oxy form (XI) can be obtained in high yield by treatment with acetic anhydride in pyridine used as a solvent. The reaction is performed at a temperature in the range of 60 - 100 °C, preferably at 80 °C, and is completed in 1 - 3 hours. This reaction may also be performed with p-toluenesulfonyl chloride or chloroform used as a solvent. Conversion from the general formula (XI) to (XII) is performed by reacting the compound of (XI) with $(\text{R}_\text{E})_3\text{OBF}_4$ (R_E is a lower alkyl group such as methyl or ethyl) or $\text{AgBF}_4 \cdot \text{R}_\text{E}\text{X}$ (R_E has the same meaning as defined above) under cooling with ice in a nitrogen stream. The reaction is performed over 4 - 7 days in a solvent such as chloroform or methylene chloride. Conversion from the general formula (XII) to (XIII) and from (XIII) to (XIV) is performed by reduction reaction in the same manner as already described.

(f) If R₁ is a hydrogen atom, a linear lower alkyl group or a hydroxyl-substituted linear lower alkyl group, the following scheme may be employed:




(where R₂, R₃, R_E, R_B and X have the same meanings as defined above).

The conversion from the general formula (II) to (IV) is carried out as in (c). Conversion from the general

formula (IV) to (XV) is accomplished by reduction reaction which is performed in the presence of a catalyst such as Raney nickel in a solvent such as tetrahydrofuran or an alcohol, preferably at room temperature. The reaction may also proceed in the presence of an iron catalyst in a solvent such as acetic acid or trifluoroacetic acid at a temperature of 90 - 100° C. If desired, further reduction may be performed with a suitable catalyst such as platinum dioxide, thereby producing an end compound of the general formula (XVI).


(g) If R_1 is a hydroxyl group, the following scheme may be employed:



25 (where R_2 , R_3 , R_A , R_B and X have the same meanings as defined above).

The conversion from the general formula (II) to (IV) and from (IV) to (XI) is performed as described above. Conversion from the general formula (XI) to (XVII) is performed as already-described in connection with the reaction for reducing the nitro group and may be accomplished by reduction in the presence of a catalyst such as palladium-on-carbon or Raney nickel. If desired, further reduction may be performed to obtain an end compound represented by the general formula (XVIII).

(h) If R_1 is a morpholino group, the following scheme may be employed:



(where R_2 , R_3 , R_A , R_B and X have the same meanings as defined above).

50 The conversion from the general formula (II) to (IV) is performed as in (c). Conversion from the general formula (IV) to (XIX) is performed by reaction with an enamine having a morpholino group such as 1-(N-morpholino)cyclohexene and an acylating agent such as benzoyl chloride, tosyl chloride or an acid anhydride such as acetic anhydride in an inert solvent such as chloroform or methylene chloride. The reaction temperature is in the range of 0 - 10° C and the reaction time is in the range of 3 - 24 hours, preferably 12 hours. The enamine is added in an amount of 1.5 - 3 equivalents, preferably 2 equivalents, and the acylating agent is added in an amount of 0.5 - 2 equivalents, preferably 1 equivalent.

55 Conversion from the general formula (XIX) to (XX) involves the usual practice of reduction as described above and may be accomplished by hydrogenation in the presence of a catalyst such as palladium-on-

carbon, Raney nickel or Raney cobalt.

The compounds of the general formula (I) thus obtained exhibit excellent activity in accelerating mnemonic and learning performance and improving the brain function in small doses and hence are useful as pharmaceutical drugs.

5 The compounds of the general formula (I) may be converted to acid addition salts as required. If such acid addition salts are to be used in medical applications, any pharmaceutically acceptable salt forming acids may be used. Specific examples include organic acids such as citric acid, fumaric acid, maleic acid and tartaric acid, and inorganic salts such as hydrochloric acid, hydrobromic acid, nitric acid and sulfuric acid.

10 The following examples and test examples are provided for the purpose of further illustrating the present invention but are in no way to be taken as limiting.

Example 1

15

Preparation of 4-amino-3-morpholino-2-(2'-oxocyclohexyl)quinoline (Compound No. 3):

20 (a) 3-Bromo-4-nitroquinoline-N-oxide (2.69 g, 0.01 mol) and 1-(N-morpholino)cyclohexene (4.18 g, 0.025 mol) were dissolved in 50 ml of chloroform and the mixture was stirred at room temperature for 4 days. After distilling off the chloroform under vacuum, the residue was subjected to extraction with benzene. Following washing with water and drying, the benzene was distilled off under vacuum and ethanol was added to the residue, whereupon crystal was precipitated. The resulting crystal was recrystallized from ethanol to obtain 3-morpholino-4-nitro-2-(2'-oxocyclohexyl)quinoline in an amount of 2.27 g (yield, 64%).

25 (b) 3-Morpholino-4-nitro-2-(2'-oxocyclohexyl)quinoline (2.00 g, 0.006 mol) was dissolved in tetrahydrofuran (10 ml) and subjected to reduction in the presence of 10% palladium-on-carbon (0.3 g). After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from ethanol was performed, whereupon 4-amino-3-morpholino-2-(2'-oxocyclohexyl)quinoline was obtained in an amount of 0.94 g (yield, 46%).

30

Example 2

35 Preparation of 4-amino-2-(2'-hydroxycyclohexyl)-3-morpholino-5,6,7,8-tetrahydroquinoline (Compound No. 6):

40 The 4-amino-3-morpholino-2-(2'-oxocyclohexyl)quinoline (0.8 g, 0.0024 mol) obtained in Example 1-(b) was dissolved in 10 ml of trifluoroacetic acid and subjected to reduction in the presence of platinum (IV) dioxide (0.3 g). After completion of the reaction, the platinum dioxide was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from benzene/n-hexane was performed, whereupon 4-amino-2-(2'-hydroxycyclohexyl)-3-morpholino-5,6,7,8-tetrahydroquinoline was obtained in an amount of 0.69 g (yield, 85%).

45

Example 3

50 Preparation of 4-amino-2-cyclohexyl-3-morpholinoquinoline (Compound No 18):

(a) The 4-amino-3-morpholino-2-(2'-oxocyclohexyl)quinoline (1.0 g, 0.0031 mol) obtained in Example 1-(b) and p-toluenesulfonyl hydrazide (0.72 g, 0.0039 mol) were dissolved in 10 ml of methanol and the solution was stirred at room temperature for 4 hours, whereupon gradual crystallization occurred. The crystal was recovered by filtration under vacuum and washed with methanol to obtain 2-(4-amino-3-morpholinoquinolyl)-1-tosylhydrazonocyclohexane in an amount of 0.65 g (yield, 43%).

55

(b) The 2-(4-amino-3-morpholinoquinolyl)-1-tosylhydrazonocyclohexane (0.65 g, 0.0013 mol) was suspended in methanol (30 ml) and 1.0g of sodium borohydride was gradually added under cooling with

ice. Thereafter, the mixture was heated under reflux on an oil bath for 5 hours. Methanol was distilled off from the whole reaction mixture under vacuum and the residue was subjected to extraction with chloroform. The chloroform layer was washed with water, dried with anhydrous sodium sulfate and subjected to isolation and purification steps on an alumina column, thereby obtaining 4-amino-2-cyclohexyl-3-morpholinoquinoline in an amount of 0.31 g (yield, 78%).

Example 4

10 Preparation of 4-amino-2-cyclohexyl-3-morpholino-5,6,7,8-tetrahydroquinoline (Compound No. 19):

The 4-amino-cyclohexyl-3-morpholinoquinoline (0.31 g, 0.001 mol) obtained in Example 3-(b) was dissolved in 10 ml of trifluoroacetic acid and subjected to reduction in the presence of platinum (IV) dioxide (0.3 g). After completion of the reaction, the platinum dioxide was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from benzene/n-hexane was performed, whereupon 4-amino-2-cyclohexyl-3-morpholino-5,6,7,8-tetrahydroquinoline was obtained in an amount of 0.25 g (yield, 83%).

Example 5

Preparation of 4-amino-2-anilino-3-morpholinoquinoline (Compound No. 13):

25 (a) 3-Bromo-4-nitroquinoline-N-oxide (10 g, 0.037 mol) was dissolved in tetrahydrofuran and after adding morpholine (8.1 g, 0.093 mol), the mixture was stirred at room temperature, whereupon gradual crystallization occurred. Tetrahydrofuran was distilled off from the whole reaction mixture under vacuum and the residue was dissolved in chloroform. The chloroform layer was washed with water and dried with anhydrous sodium sulfate. After distilling off the chloroform under vacuum, the residue was washed with a small amount of ethanol and recovered by filtration under vacuum, whereupon 3-morpholino-4-nitroquinoline-N-oxide was obtained in an amount of 10 g (yield, 98%).

(b) The 3-morpholino-4-nitroquinoline-N-oxide (3 g, 0.011 mol) was suspended in dimethylformamide (40 ml) and following addition of phenyl isocyanate (3.4 g, 0.029 mol), the mixture was heated on an oil bath at 80 - 90 °C for ca. 3 hours. The reaction mixture was totally charged into water and subjected to extraction with chloroform. The chloroform layer was washed with water and dried with anhydrous sodium sulfate. After the chloroform was distilled off under vacuum, the residue was subjected to recrystallization with ethanol. The resulting crystal was recovered by filtration under vacuum to obtain 2-anilino-3-morpholino-4-nitroquinoline in an amount of 2.95 g (yield, 77%).

40 (c) The 2-anilino-3-morpholino-4-nitroquinoline (2.9 g, 0.0083 mol) was dissolved in tetrahydrofuran (10 ml) and subjected to reduction in the presence of 10% palladium-on-carbon (0.3 g). After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from methanol was conducted to obtain 4-amino-2-anilino-3-morpholinoquinoline in an amount of 1.89 g (yield, 69%).

Example 6

Preparation of 4-amino-2-anilino-3-morpholino-5,6,7,8-tetrahydroquinoline (Compound No. 15):

50 The 4-amino-2-anilino-3-morpholinoquinoline (1.8 g, 0.0055 mol) obtained in Example 5-(c) was dissolved in trifluoroacetic acid (10 ml) and subjected to reduction in the presence of platinum (IV) dioxide (0.3 g). After completion of the reaction, platinum dioxide was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from benzene was performed, whereupon 4-amino-2-anilino-3-morpholino-5,6,7,8-tetrahydroquinoline was obtained in an amount of 0.58 g (yield, 32%).

Example 7

Preparation of 4-amino-2-ethylamino-3-morpholinoquinoline (Compound No. 43):

5

(a) The 3-morpholino-4-nitroquinoline-N-oxide (3 g, 0.011 mol) obtained in Example 5-(a) was subjected in dimethylformamide (40 ml) and following the addition of ethyl isocyanate (2.7 g, 0.038 mol), the suspension was heated in a sealed tube on an oil bath at 130 °C for ca. 6 hours. The mixture was totally charged into water and subjected to extraction with chloroform. The chloroform layer was washed with water, dried with anhydrous sodium sulfate and concentrated under vacuum. Following isolation by chromatography on a silica gel column (solvent: 5% methanol/chloroform), recrystallization from ethanol was performed, whereupon 2-ethylamino-3-morpholino-4-nitroquinoline was obtained in an amount of 2.1 g (yield, 64%).

10

(b) The 2-ethylamino-3-morpholino-4-nitroquinoline (2.0 g, 0.0066 mol) was dissolved in tetrahydrofuran (10 ml) and subjected to reduction in the presence of 10% palladium-on-carbon (0.3 g). After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from ethanol was conducted, whereupon 4-amino-2-ethylamino-3-morpholinoquinoline was obtained in an amount of 1.2 g (yield, 67%).

20

Example 8

Preparation of 4-amino-2-ethoxy-3-morpholinoquinoline (Compound No. 38):

25

(a) The 3-morpholino-4-nitroquinoline-N-oxide (4 g, 0.015 mol) obtained in Example 5-(b) was dissolved in acetic anhydride (20 ml) and pyridine (20 ml) and the solution was heated on an oil bath at 80 °C for 3 hours. Excess acetic anhydride and pyridine were distilled off from the whole reaction mixture in vacuum and the residue was washed with ether to obtain 3-morpholino-4-nitro-2-quinolone in an amount of 3.5 g (yield, 88%).

30

(b) The 3-morpholino-4-nitro-2-quinolone (3.5 g, 0.013 mol) was suspended in methylene chloride (200 ml). To the ice-cooled suspension, 25 ml of a solution of triethyl oxonium tetrafluoroborate in methylene chloride was added in a nitrogen stream and the mixture was stirred at room temperature for 7 days. The whole portion of the mixture was poured into 50% aqueous potassium carbonate and the resulting mixture was subjected to extraction with chloroform. The chloroform layer was washed with water and dried with anhydrous sodium sulfate. Upon isolation and purification by chromatography on a silica gel column (solvent: chloroform), 2-ethoxy-3-morpholino-4-nitroquinoline was obtained in an amount of 1.3 g (yield, 34%).

35

(c) The 2-ethoxy-3-morpholino-4-nitroquinoline (1.2 g, 0.004 mol) was dissolved in tetrahydrofuran (10 ml) and subjected to reduction in the presence of 10% palladium-on-carbon (0.3 g). After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from chloroform was performed to obtain 4-amino-2-ethoxy-3-morpholinoquinoline in an amount of 0.7 g (yield, 60%).

45

Example 9

Preparation of 4-amino-3-piperidino-2-(2'-oxocyclohexyl)quinoline (Compound No. 7):

50

(a) 3-Bromo-4-nitroquinoline-N-oxide (10 g, 0.037 mol) and 1-(N-piperidino)cyclohexene (15.4 g, 0.093 mol) were dissolved in chloroform (100 ml) and the solution was stirred at room temperature for 2 days. The chloroform was distilled off under vacuum and the residue was subjected to extraction with benzene. Following washing with water and drying, benzene was distilled off under vacuum and the residue was purified and isolated by chromatography on a silica gel column using chloroform as a developing solvent. By recrystallization from ethanol, 3-piperidino-4-nitro-2-(2'-oxocyclohexyl)quinoline was obtained in an amount of 9 g (yield, 68%).

55

(b) The 3-piperidino-4-nitro-2-(2'-oxocyclohexyl)quinoline (8.5 g, 0.024 mol) was dissolved in

tetrahydrofuran (50 ml) and subjected to reduction in the presence of 10% palladium-on-carbon (1.2 g). After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from ethanol was conducted to obtain 4-amino-3-piperidino-2-(2'-oxocyclohexyl)quinoline in an amount of 6.1 g (yield, 74%).

Example 10

10 Preparation of 4-amino-2-hydroxy-3-morpholinoquinoline (Compound No. 27):

(a) The 3-morpholino-4-nitroquinoline-N-oxide obtained in Example 5-(a) was dissolved in acetic anhydride (16 ml) and pyridine (16 ml) and the solution was heated on an oil bath at 80 °C for 1 hour. Excess acetic anhydride and pyridine were distilled off under vacuum from the whole reaction mixture and the residue was dissolved in chloroform, followed by washing with 10% NaHCO₃. The insoluble matter was recovered by filtration under vacuum, dissolved in acetone and treated with charcoal. The chloroform layer was dried with anhydrous Na₂SO₄ and CHCl₃ was distilled off under vacuum. The residue was combined with the charcoal treated (insoluble) matter and washed with ethanol to obtain 2-hydroxy-3-morpholino-4-nitroquinoline in an amount of 3.5 g (yield, 44%).

20 (b) The 2-hydroxy-3-morpholino-4-nitroquinoline (1.5 g, 0.0054 mol) was dissolved in a mixed solvent of methanol and tetrahydrofuran and subjected to catalytic reduction in the presence of 10% palladium-on-carbon. After completion of the reaction, the palladium-on-carbon catalyst was filtered off and the mother liquor was concentrated. Following purification by chromatography on an alumina column using chloroform as a developing solvent, recrystallization from chloroform was conducted to obtain 4-amino-2-hydroxy-3-morpholinoquinoline in an amount of 0.9 g (yield, 63%).

Example 11

30 Preparation of 4-amino-3-piperidinoquinoline (Compound No. 1):

(a) 3-Bromo-4-nitroquinoline-N-oxide (10 g, 0.037 mol) was dissolved in tetrahydrofuran (100 ml). To the solution, piperidine (7.9 g, 0.093 mol) was added and the mixture was stirred at room temperature until gradual crystallization occurred. Tetrahydrofuran was distilled off from the whole reaction mixture under vacuum and the residue was dissolved in chloroform. The chloroform layer was washed with water and dried with anhydrous sodium sulfate. After distilling off the chloroform under vacuum, the residue was washed with a small amount of ethanol and recovered by filtration under vacuum to obtain 4-nitro-3-piperidinoquinoline-N-oxide in an amount of 6.0 g (yield, 59%).

40 (b) The 4-nitro-3-piperidinoquinoline-N-oxide (4 g, 0.015 mol) was dissolved in a mixed solvent of methanol and tetrahydrofuran (50 ml) and subjected to catalytic reduction in the presence of purified Raney nickel (2 ml). Upon isolation and purification by chromatography on an alumina column (developing solvent: CHCl₃), 4-amino-3-piperidinoquinoline was obtained in an amount of 2.1 g (yield, 63%).

Example 12

45

Preparation of 4-amino-3-ethylamino-2-morpholinoquinoline (Compound No. 52):

50 (a) 3-Bromo-4-nitroquinoline-N-oxide (10 g, 0.037 mol) was dissolved in tetrahydrofuran (100 ml). To the solution, ethylamine (5.96 g, 70% in H₂O) was added and the mixture was stirred at room temperature for 1 hour. Tetrahydrofuran was distilled off under vacuum from the whole reaction mixture and the residue was dissolved in chloroform. The chloroform layer was washed with water and dried with anhydrous sodium sulfate. Upon isolation and purification by chromatography on a silica gel column using chloroform as a developing solvent, 3-ethylamino-4-nitroquinoline-N-oxide was obtained in an amount of 5.4 g (yield, 62%).

55 (b) The 3-ethylamino-4-nitroquinoline-N-oxide (1 g, 0.004 mol) was dissolved in chloroform (15 ml). To the ice-cooled solution, 1-(N-morpholino)-cyclohexene (1.7 g) and benzoyl chloride (0.7 g) were added dropwise. The mixture was restored to room temperature and stirred overnight. Following addition of water, the mixture was stirred for about 1 hour and washed with water. Thereafter, the CHCl₃ layer was separated

and dried with anhydrous Na_2SO_4 . Upon isolation and purification by chromatography on a silica gel column using chloroform as a developing solvent, 3-ethylamino-2-morpholino-4-nitroquinoline was obtained in an amount of 0.7 g (yield, 54%).

5 (c) The 3-ethylamino-2-morpholino-4-nitroquinoline (0.7 g, 0.0023 mol) was dissolved in a mixed solvent of methanol and tetrahydrofuran (15 ml) and subjected to catalytic reduction in the presence of 10% palladium-on-carbon (0.3 g), thereby obtaining 4-amino-3-ethylamino-2-morpholinoquinoline in an amount of 0.5 g (yield, 79%).

10 The compounds prepared in Examples 1 - 12 and other compounds of the general formula (I) are characterized in Table 1 below, wherein "Q" in the " R_4 , R_5 " column means that R_4 and R_5 taken together with the ring A form a quinoline ring; "T" means that R_4 and R_5 taken together with the ring A form a 5,6,7,8-tetrahydroquinoline ring; and "P" means that R_4 and R_5 each represents a hydrogen atom.

15

20

25

30

35


40

45

50

55

Table 1



Comp. No.	R ₁	R ₂	R ₃	R ₄ · R ₅	m.p. (°C)	NMR	salt
1	H			Q	255 - 257	1.36-2.27(6H,m), 2.72-3.26(4H,m), 5.32(2H,bs), 7.11-8.03(4H,m), 8.56(1H,s)	2HCl
2	H			Q	113 - 114	2.32(3H,s), 2.35-3.20(8H,m), 5.30(2H,bs), 7.11-8.12(4H,m), 8.49(1H,s)	
3				Q	195 - 196	1.50-2.87(8H,m), 3.01(4H,m), 3.79(4H,m), 4.12(1H,dd), 5.10(2H,bs), 7.15-8.05(4H,m)	
4	-CH ₂ OH	H	H	Q	242 - 244	3.41(1H,b), 4.85(2H,b), 5.11(2H,bs), 7.04-8.49(4H,m), 8.06(2H,bs)	
5	-CH ₂ OH	H	COCH ₃	Q	218 - 224 (dec.)	2.08(3H,s), 3.26(1H,b), 4.46(2H,bs), 6.31(2H,bs), 7.05-8.30(4H,m), 8.86(1H,bs)	
6				T	247 - 248	1.45-3.37(21H,m), 3.63-3.95(4H,m), 4.09(1H,s), 4.79(2H,bs), 7.50(1H,bs)	

Table 1 (cont'd)













Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
7				Q	192 - 194	1.50-1.90(6H,m), 1.90-2.80(8H,m), 2.80-3.25(4H,m), 4.12(1H,dd), 5.12(2H,bs), 7.15-8.05(4H,m)	
8	-CH ₃			Q	270 - 280 (sublimable)	2.87(3H,s), 2.90-4.44(8H,m), 5.10(2H,bs), 7.49-8.31(4H,m)	
9				T	220 - 223	1.32-3.00(27H,m), 4.08(1H,s), 4.72(2H,bs), 7.56(1H,bs)	
10	-CH ₃			T	133 - 134	1.65-2.88(8H,m), 2.42(3H,s), 3.10-4.21(8H,m), 4.67(2H,bs)	
11				Q	205 - 208	0.98(3H,t), 1.51(3H,d), 2.17-2.59(2H,q), 2.66-3.98(8H,m), 3.92-4.39(1H,q), 5.35(2H,bs), 7.13-8.01(4H,m)	
12				Q	-	-	
13				Q	224 - 225	2.51-3.79(4H,m), 3.70-4.11(4H,m), 5.59(2H,bs), 6.70-7.97(9H,m), 8.51(1H,bs)	

Table 1 (cont'd)












Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
14	H			Q	168 - 170	2.21-3.15(4H,m), 3.16-4.07(4H,m), 5.30(2H,bs), 7.16-8.09(4H,m), 8.56(1H,s)	
15	 -NH-			T	184 - 186	1.62-2.86(8H,m), 2.90-3.62(4H,m), 3.63-3.99(4H,m), 4.08(2H,bs), 6.66-8.01(6H,m)	
16	 -NH-			T	189 - 191	1.02-3.46(22H,m), 3.50-4.24(5H,m), 5.74(2H,bs), 6.69(1H,bs)	
17	H			T	136 - 138	1.67-3.20(12H,m), 3.70-4.05(4H,m), 4.43(2H,bs), 7.92(1H,s)	
18				Q	195 - 196	1.11-2.12(10H,m), 2.53-4.25(9H,m), 5.30(2H,bs), 7.12-8.02(4H,m)	
19				T	191 - 193	1.10-3.07(22H,m), 3.10-4.11(5H,m), 4.59(2H,bs)	
20	H			Q	263 - 264 (dec.)	2.04-2.37(4H,m), 3.82-4.18(4H,m), 5.01(2H,bs), 7.35-8.51(5H,m)	

Table 1 (cont'd)








Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
21	H	H		Q	-	-	
22		-CH ₂ -CH ₃	-CH ₂ -CH ₃	Q	192 - 193	1.02(6H, t), 1.50-2.90(8H, m), 2.80-3.31(4H, m), 5.17(2H, bs), 7.19-8.02(4H, m)	
23		-CH ₂ -CH ₃	-CH ₂ -CH ₃	T	123 - 125	1.01(6H, t), 1.20-3.38(21H, m), 4.02(1H, s), 4.70(2H, bs), 7.75(1H, bs)	
24		-CH ₂ -CH ₃	-CH ₂ -CH ₃	Q	280 - 285 (dec.)	1.03(6H, t), 1.20-3.70(15H, m), 5.23(2H, bs), 7.20-8.00(4H, m)	HCl
25		-CH ₂ -CH ₃	-CH ₂ -CH ₃	T	265 - 270 (sublimable)	1.01(6H, t), 1.10-3.35(22H, m), 3.40-3.86(1H, m), 4.57(2H, bs)	
26				Q	131 - 132	1.09-2.47(16H, m), 2.64-3.39(5H, m), 5.30(2H, bs), 7.06-7.97(4H, m)	
27	-OH			Q	283 - 284	2.91-4.20(8H, m), 6.11(2H, bs), 6.72-8.05(4H, m), 11.15(1H, bs)	

Table 1 (cont'd)









Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
28				T	144 - 146	1.01-3.45(28H,m), 4.20-4.51(1H,m), 5.30(2H,bs)	
29		-CH ₂ CH ₂ CH ₃	-CH ₂ CH ₂ CH ₃	Q	176 - 177	0.64-1.10(6H,m), 1.10-3.40(6H,m), 3.85-4.30(1H,dd), 5.18(2H,bs), 7.13-8.07(4H,m)	
30		-CH ₂ CH ₂ CH ₃	-CH ₂ CH ₂ CH ₃	T	123 - 124	0.52-1.10(6H,m), 1.20-3.25(26H,m), 4.11(1H,bs), 6.72(2H,bs)	
31		-CH ₂ CH ₂ CH ₃	-CH ₂ CH ₂ CH ₃	Q	129 - 130	0.67-1.03(6H,m), 1.10-2.30(14H,m), 2.60-3.35(5H,m), 5.22(2H,bs), 7.03-7.99(4H,m)	
32		-CH ₂ CH ₂ CH ₃	-CH ₂ CH ₂ CH ₃	T	119 - 120	0.71-1.05(6H,m), 1.10-2.57(22H,m), 2.60-3.23(5H,m), 4.52(2H,bs)	
33		-CH ₂ CH ₃	-CH ₂ CH ₃	Q	153 - 155	0.88-1.27(6H,t), 2.95-3.30(4H,q), 3.97(1H,bs), 4.46(2H,bs), 6.75-8.28(9H,m)	
34		-CH ₂ CH ₃	-CH ₂ CH ₃	T	224 - 226	1.04(6H,t), 1.51-2.03(4H,m), 2.12-2.84(4H,m), 2.97-3.65(4H,q), 7.20-8.35(7H,m), 12.97(1H,bs)	2HCl

Table 1 (cont'd)














Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
35	 -NH-	 -N-		Q	119 - 121	1.34-2.06(6H,m), 2.77-3.34(4H,m), 4.49(2H,bs), 6.78-8.06(9H,m); 8.33(1H,bs)	
36	 -NH-	 -N-		T	270 (dec.)	1.69-3.21(24H,m), 3.62-4.22(6H,m), 4.41-4.58(1H,m), 7.84(1H,b)	2HCl
37	 HO	 -N-		T	215 - 218	0.98(3H,t), 2.39(3H,d), 1.60-4.36(21H,m), 4.61(2H,bs)	HCl
38	-OCH ₂ CH ₃	 -N-		Q	149 - 151	1.45(3H,t), 2.32-4.09(2H,m), 4.33-4.68(2H,q), 5.19(2H,bs), 7.03-7.75(4H,m)	
39		 -N-		Q	132 - 134	0.65-2.20(10H,m), 2.61-4.22(9H,m), 5.29(2H,bs), 7.25-8.01(4H,m)	
40		 -N-		T	178 - 179	0.67-1.12(3H,m), 1.30-1.55(3H,d), 1.61-4.16(21H,m), 6.82(2H,bs)	
41		 -N-		T	137 - 140	1.54-3.33(20H,m), 3.60-4.05(5H,m), 4.47(2H,bs)	

Table 1 (cont'd)













Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
42	-OCH ₂ CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	Q	167 - 169 (dec.)	1.08(6H,t), 1.47(3H,t), 3.31(4H,q), 4.83(2H,q), 7.12-7.89(4H,m), 8.10(2H,bs)	
43	-NHCH ₂ CH ₃			Q	152 - 153	1.24(3H,t), 2.61-3.93(10H,m), 5.44(2H,bs), 5.65(1H,bs), 6.81-7.89(4H,m)	
44	-OCH ₂ CH ₃			Q	168 - 170	1.40-2.06(9H,m), 2.10-3.61(4H,m), 4.63-5.03(4H,q), 6.96-7.72(4H,m), 8.11-8.5(2H,m)	
45	-NHCH ₂ CH ₃			Q	99 - 100	1.29(3H,t), 1.30-2.18(6H,m), 2.71-3.35(4H,m), 3.30-4.11(2H,m), 4.25-4.92(2H,bs), 5.21-5.70(1H,b), 6.80-7.99(4H,m)	
46	-NHCH ₂ CH ₃			Q	99 - 101	1.23(3H,t), 1.84-2.31(4H,m), 3.02-3.38(4H,m), 3.38-3.87(2H,m), 4.44(2H,bs), 5.10(1H,m), 6.82-7.78(4H,m)	
47	-OCH ₂ CH ₃	-COCH ₃	-CH ₂ CH ₃	Q	117 - 119	1.10(3H,t), 1.34(3H,t), 1.82(3H,s), 3.43-3.89(2H,m), 4.25-4.69(2H,q), 4.99(2H,bs), 7.05-7.84(4H,m)	

Table 1 (cont'd)

Comp. No.	R ₁	R ₂	R ₃	R ₄ ·R ₅	m.p. (°C)	NMR	salt
48				Q	205 - 207	0.80-1.15(3H,m), 1.21-2.06(10H,m), 2.54-3.53(4H,m), 3.67-4.12(2H,m), 6.52-6.98(2H,bs), 6.99-8.67(4H,m)	
49				P	175 - 176	1.10-2.80(8H,m), 2.94(4H,m), 3.72(4H,m), 3.91(1H,dd), 6.34-8.00(2H,ABq)	
50				P	154 - 155	1.30-1.81(6H,m), 1.81-2.70(8H,m), 2.70-3.20(4H,m), 3.94(1H,dd) 4.51(2H,bs), 6.33-8.00(2H,ABq)	
51		-CH ₂ CH ₃	-CH ₂ CH ₃	P	146 - 147	1.00(6H,t), 1.35-2.66(8H,m), 2.86(4H,q), 3.85(1H,dd), 4.53(2H,bs), 6.31-8.04(2H,ABq)	
52		H	-CH ₂ CH ₃	Q	89 - 90	1.19(3H,t), 1.57-2.20(4H,m), 2.51-2.90(4H,m), 4.10-4.91(5H,m), 7.00-8.45(4H,m)	

(dec.) denotes decomposition point.

Test Example 1

5

Inhibition of Acetylcholinesterase Activity:

10 Acetylcholinesterase activity was assayed in accordance with the method of S.H. Sterri and F. Fonnum described in European Journal of Biochemistry, 91: 215 - 222, 1978. The method consisted of incubating 0.5 μ U of authentic acetylcholinesterase (EC3,1,1,7) (Sigma, No. C-2888) at 30 °C for 15 minutes in a 20 mM sodium phosphate buffer (pH 7.4), with 1.3 mM [14 C] acetylcholine (0.025 μ Ci) being used as a substrate. Selected compounds of the general formula (I) were added at a concentration of 1 μ M to the reaction solution and the resulting effect on enzymatic activity was evaluated.

15 The result was obtained as the average of three measurements and shown in Table 2 in terms of percentage, with the activity for the case where no drum compound was added being taken as 100.

20

25

30

35

40

45

50

55

Table 2













Compound No.	Structure	Concentration	Inhibition rate (%)
None	-	-	100
2		1 μ M	27.1
6		1 μ M	36.9
7		1 μ M	38.1
9		1 μ M	38.8
15		1 μ M	37.2
16		1 μ M	25.9
19		1 μ M	21.8
25		1 μ M	30.1

Table 2 (cont'd)

Compound No.	Structure	Concentration	Inhibition rate (%)
28		1 μ M	22.2
32		1 μ M	28.7
35		1 μ M	23.3
41		1 μ M	4.2

As Table 2 shows, the tested compounds of the present invention have a strong power to inhibit acetylcholinesterase activity and hence are expected to stimulate the cholinergic function of the central nervous system. This will justify the anticipation that the compounds of the general formula (I) according to the present invention will be useful in the treatment of dementia of Alzheimer type which is accompanied by an observable damage specific to central cholinergic neurons (P.J. Whitehouse et al., Science, 215, 1237 - 1239, 1982).

Test Example 2

Inhibition of Catecholamine Uptake:

A crude synaptosomal fraction (P2) was obtained from the brain of rats (F344/C₇₁) in the usual manner. In accordance with the method of S.H. Snyder and J.T. Coyle described in Journal of Pharmacology and Experimental Therapeutics, 165(1), 78 - 86, 1969, the fraction P2 was incubated in a Krebs-Henseleit bicarbonate buffer solution at 37°C for 10 minutes together with 50 nM ³H-norepinephrine (0.25 μ Ci). Thereafter, the uptake of norepinephrine into the synaptosomes was measured by the filtration method. Selected compounds were added at a concentration of 1 μ M to the solution being incubated and their effect on norepinephrine uptake was examined. The control drug was tetrahydroaminoacridine (THA).

The results are shown in Table 3 in terms of percentage, with norepinephrine uptake in the absence of drug compounds being taken as 100.

Table 3











Compound No.	Structure	Concentration	Inhibition rate (%)
None	-	-	100
1		1 μ M	69.1
6		1 μ M	82.6
7		1 μ M	79.0
9		1 μ M	69.4
15		1 μ M	85.4
16		1 μ M	88.1
19		1 μ M	83.2
23		1 μ M	73.7

Table 3 (cont'd)

Compound No.	Structure	Concentration	Inhibition rate (%)
24		1 μ M	68.2
25		1 μ M	75.3
26		1 μ M	68.2
28		1 μ M	85.2
30		1 μ M	76.9
32		1 μ M	73.2
34		1 μ M	80.0
39		1 μ M	71.9
43		1 μ M	78.8
45		1 μ M	79.8

Table 3 (cont'd)

Compound No.	Structure	Concentration	Inhibition rate (%)
52		1 μ M	79.8
control		1 μ M	84.6

As Table 3 shows, in most of the cases examined, the tested compounds of the present invention exhibited a stronger activity than THA which is known as an effective drug against senile dementia. This result shows that the compounds of the present invention have a strong power to stimulate catecholaminergic neurons in the central nervous system, thus clearly demonstrating their utility as drugs in the treatment of neural and mental disorders.

Test Example 3

Improvement of memory and Learning Ability Tested by Passive Avoidance Response:

Four groups of male ICR mice (10 - 14 weeks old), each consisting of 10 animals, were used in the experiment. Except for one control group, the mice were administered with selected compounds of the present invention. The experimental chamber consisted of a light and a dark compartment and an electric current shock source was connected to the floor grid in the dark compartment.

The experimental sessions consisted of acclimation, acquisition, and retention trials and were performed for 3 days, one trial per day. In the acclimation trial, animals were placed in the light compartment of the experimental chamber and left there for 5 minutes to acclimate them to the apparatus. In the acquisition trial which was conducted on the 2nd day, the animals were placed in the light compartment and when they entered the dark compartment, they were confined and given an a.c. electric shock (1 mA) through the floor grid for 10 seconds. In the retention trial which was conducted 24 hours after the acquisition trial, the animals were placed in the light compartment of the chamber and the step-through latency (i.e., the time required for the animals to enter the dark compartment) was measured. A physiological saline solution was administered at the acclimation trial to all of the animals. At the acquisition trial, selected compounds were administered in amounts of 10 μ g/kg of body weight (1 ml/100 g). The control group was administered the same amount of physiological saline solution 60 minutes before the start of the trial. All administrations were intraperitoneal.

The results are shown in Table 4 below.

Table 4









Compound No.	Structure	dose ($\mu\text{g/kg i.p.}$)	Step through latency (sec)
None	-	-	41.9 ± 10.2
2		10	92.1 ± 13.5
6		10	144.8 ± 45.1
7		10	104.1 ± 28.5
9		10	54.4 ± 7.2
13		10	100.2 ± 17.8
15		10	80.4 ± 14.6
16		10	102.1 ± 22.3
18		10	116.4 ± 29.2

Table 4 (cont'd)

Compound No.	Structure	dose (µg/kg i.p.)	Step through latency (sec)
19		10	107.1 ± 29.9
22		10	144.8 ± 25.5
24		10	94.2 ± 14.7
32		10	109.7 ± 16.6
39		10	108.3 ± 16.0
40		10	113.1 ± 29.6
43		10	109.1 ± 13.4
44		10	90.4 ± 12.4
45		10	70.7 ± 6.8
50		10	83.4 ± 14.8

As Table 4 shows, the step-through latency of the control group which was 41.9 ± 10.2 was significantly delayed by the tested compounds of the present invention, which hence were found to be effective in promoting memory and learning performance.

5

Test Example 4

10 Inhibition of Catecholamine Uptake:

The uptake of dopamine into synaptosomes was measured by repeating the procedure of Test Example 2 except that 50 nM ^3H -norepinephrine (0.25 μCi) was replaced by 15 nM ^3H -dopamine (0.68 μCi). The control drug was tetrahydroaminoacridine (THA). The result is shown in Table 5 in terms of percentage, with
15 dopamine uptake in the absence of drugs being taken as 100.

20

25

30

35

40

45






50

55

Table 5

Compound No.	Structure	Concentration	Inhibition rate (%)
None	-	-	100
7		1 μ M	86.4
9		1 μ M	78.6
23		1 μ M	83.0
24		1 μ M	59.6
25		1 μ M	76.4
26		1 μ M	55.4
28		1 μ M	80.5
30		1 μ M	79.2
32		1 μ M	71.0

Table 5 (cont'd)

Compound No.	Structure	Concentration	Inhibition rate (%)
34		1 μ M	84.8
45		1 μ M	82.3
49		1 μ M	67.5
52		1 μ M	57.8
control		1 μ M	91.5







As Table 5 shows, the tested compounds of the present invention had a stronger activity than THA, indicating together with the results shown in Test Example 2 their great ability to stimulate catecholaminergic neurons in the central nervous system. The compounds of the present invention are therefore anticipated to prove useful as a drug in the treatment of neural and mental disorders.

Test Example 5

Inhibition of Monoamine oxidase (MAO):

A crude synaptosomal fraction (P2) was obtained from the brain of rats (F344/C₇₁) in the usual manner and homogenized to perform MAO activity measurements. The substrate was 12 μ M [¹⁴C]-serotonin in the case of MAO-A activity measurements, and 5 μ M [¹⁴C]-2-phenethylamine in the case of MAO-B activity measurements. In accordance with the method of R.J. Wurtman and J. Axelrod described in Biochemical Pharmacol. 12, 1439 - 1440, 1963, the homogenates were incubated in 100 mM phosphate buffer (pH 7.4) at 37° C for 20 minutes and the effect of adding selected compounds at a concentration of 10 μ M was evaluated. The results are shown in Table 6.

Table 6

Compound No.	Structure	Concentration	Inhibition rate (%)	
			MAO-A	MAO-B
None	-	-	100	100
1		10 μ M	44.1	82.9
7		10 μ M	58.7	87.5
9		10 μ M	59.4	89.4
49		10 μ M	54.5	71.2
52		10 μ M	0.0	63.0
control		10 μ M	60.8	90.3

As Table 6 shows, the tested compounds of the present invention were capable of inhibiting both MAO-A and MAO-B, with their ability to inhibit MAO-A being particularly great. Therefore, the compounds of the present invention are anticipated to exhibit stimulatory effect on monoaminergic neurons in the central nervous system, thereby proving useful in the treatment of neural and mental disorders caused by the functional impairment of such neurons.

Test Example 6

Effect on Performance in a Radial-Arm Maze Task:

In accordance with the method described by Thomas, J.W. et al. in Brain Research, 321, 91 - 102, 1984, 0.05 nmol of AF64A picryl sulfonate (hereunder abbreviated as AF64AP) was injected into the both sides lateral cerebral ventricles of male rats (F344) that had been made to acquire the ability to solve a radial-arm maze problem in accordance with the method of Olton and Samuelson (Olton, D.S. and Samuelson, R.J., J. Exp. Psychol., 2, 97 - 116, 1976), so as to impart a damage specific to the hippocampal


cholinergic system. The mnemonic ability of the rats was evaluated with the number of correct responses (i.e., the number of different arms selected in the first 8 choices) being used as a measure of their performance. The rats treated with AF64AP were found to have suffered significant memory impairment compared to the control group.

5 The treated animals were administered orally a vehicle on day 1, 3 and 5, Compound No. 7 on day 2, 4 and 6 (the dose was 3, 10 and 30 mg/kg on day 2, 4 and 6, respectively), and thereafter checked for their ability to solve a radial-arm maze problem. On each day when the performance subsequent to the administration of Compound No. 7 was compared with the control data (obtained by vehicle administration on the previous day) by a paired t-test, a statistically significant increase in the number of correct responses
10 was observed at 10 mg/kg ($p < 0.05$; $n = 6$) and 30 mg/kg ($p < 0.05$; $n = 6$). The results are shown in Fig. 1, with the control data for the first and last days of the test being designated by V. Chance level of correct responses defined by Spetch, M.L. and Wilkie, D.M., Behavior Research Methods and Instrumentations, 12, 377 - 378, 1980, was expressed by dashed line. All the control data obtained in the test were stable since there was no significant difference among the control data.

15 The results described above and depicted in Fig. 1 show that the compounds of the present invention will prove effective in improving the cognitive ability of patients suffering from Alzheimer's disease and other types of dementia which are characterized by extensive damage to the central cholinergic system.

20 Claims

1. A 4-aminopyridine derivative represented by the general formula I:



30 [where R_1 is a hydrogen atom, a hydroxyl group, a linear or branched alkyl group having 1 to 6 carbon atoms or a cycloalkyl group having 1 to 6 carbon atoms which may be substituted by a hydroxyl group, an alkoxy group having 1 to 6 carbon atoms, an alkyl having 1 to 6 carbon atoms or a cycloalkyl group having 1 to 6 carbon atoms which contains a carbonyl group, a morpholino group or a group $-NH-B$ (where B is an alkyl group having 1 to 6 carbon atoms, a cycloalkyl group having 1 to 6 carbon atoms or a phenyl group);
35 R_2 and R_3 which may be the same or different each represents a hydrogen atom, an alkyl group having 1 to 6 carbon atoms or an alkylcarbonyl group having 2 to 7 carbon atoms, or when taken together, form an azacycloalkyl group, a morpholino group or an N-methylpiperazinyl group together with the nitrogen atom; R_4 and R_5 each represents a hydrogen atom, or when taken together with the ring A, form a quinoline ring
40 or a 5,6,7,8-tetrahydroquinoline ring, provided that when each of R_1 , R_4 and R_5 is a hydrogen atom, R_2 and R_3 are neither a hydrogen atom nor a methyl group at the same time, and when R_1 is a hydrogen atom and R_4 and R_5 taken together with the ring A form a quinoline ring, R_2 and R_3 are neither a hydrogen atom nor an ethyl group at the same time] or an acid addition salts of said 4-aminopyridine derivative.

2. A compound according to Claim 1 wherein the cycloalkyl group has 3 - 7 carbon atoms.

45 3. A compound according to Claim 1 wherein the alkyl group is a methyl, ethyl, n-propyl, n-butyl or 2-pentyl group, the alkoxy group is a methoxy or ethoxy group, and the cycloalkyl group is a cyclohexyl group.

4. A compound according to Claim 1 wherein the alkylcarbonyl group is an acetyl or propionyl group.

5. A compound according to Claim 1 wherein the azacycloalkyl group has 3 - 6 carbon atoms.

50 6. A compound according to Claim 5 wherein the azacycloalkyl group is a pyrrolidyl or piperidyl group.

7. A compound according to Claim 1 wherein R_2 and R_3 taken together with the nitrogen atom form an azacycloalkyl, morpholino or N-methylpiperazyl group.

8. A pharmaceutical composition for improving psychoneural function which contains a pharmaceutically acceptable vehicle and a compound of the general formula according to any one of claims 1 to 7.

55 9. A pharmaceutical composition according to Claim 8 which is effective in the treatment of Alzheimer type dementia.

10. A pharmaceutical composition according to Claim 8 which is effective in the treatment of pathergasia due to impaired monoaminergic neurons.

11. A pharmaceutical composition according to Claim 8 which is effective in promoting mnemonic and learning performance.

5

10

15

20

25

30

35


40

45

50

55

Fig. 1



* $p < 0.05$, PAIRED T-TEST



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 361 489 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89118014.3

(51) Int. Cl.⁵: C07D 215/42, A61K 31/47,
C07D 213/74, A61K 31/44

(22) Date of filing: 28.09.89

(30) Priority: 30.09.88 JP 246237/88

(43) Date of publication of application:
04.04.90 Bulletin 90/14

(54) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

(68) Date of deferred publication of the search report:
12.06.91 Bulletin 91/24

(71) Applicant: Chugai Seryaku Kabushiki Kaisha
5-1, 5-chome, Ukima Kita-ku
Tokyo(JP)

(72) Inventor: Miura, Yutaka
Nochi Mansions B-201 10-20, Kobara-cho
1-chome

Mishima-shi Shizuoka-ken(JP)

Inventor: Yoshida, Mitsutaka

24-2, Fujimidai

Mishima-shi Shizuoka-ken(JP)

Inventor: Fujimura, Yasuo

2-94-8, Sakae, Ina-cho

Kitaadachi-gun Saitama-ken(JP)

Inventor: Takaku, Sakae

381-12, Oazakoizumi

Ageo-shi Saitama-ken(JP)

Inventor: Noda, Yukifumi


888-24, Shmotogari Nagaizumi-cho

Sunto-gun Shizuoka-ken(JP)

(74) Representative: Vossius & Partner
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)

(54) Novel 3,4-diaminoquinoline and pyridine compounds.

(57) 4-aminopyridine derivatives represented by the general formula:



[where R₁ is a hydrogen atom, a hydroxyl group, a linear or branched lower alkyl group or a cycloalkyl group which may be substituted by a hydroxyl group, a lower alkoxy group, a lower alkyl or a cycloalkyl group which contains a carbonyl group, a morpholino group or a group -NH-B (where B is a lower alkyl group, a cycloalkyl group or a phenyl group); R₂ and R₃ which may be the same or different each represents a hydrogen atom, a lower alkyl group or a loweralkylcarbonyl group, or when taken together, form an azacycloalkyl group, a morpholino group or an N-methylpiperazinyl group to-

gether with the nitrogen atom; R₄ and R₅ each represents a hydrogen atom, or when taken together with the ring A, form a quinoline ring or a 5,6,7,8-tetrahydroquinoline ring, provided that when each of R₁, R₄ and R₅ is a hydrogen atom, R₂ and R₃ are neither a hydrogen atom nor a methyl group at the same time, and when R₁ is a hydrogen atom and R₄ and R₅ taken together with the ring A form a quinoline ring, R₂ and R₃ are neither a hydrogen atom nor an ethyl group at the same time], and an acid addition salts of said 4-aminopyridine derivatives. These compounds are useful as active ingredients in pharmaceutical compositions for improving psychoneural function, especially in the treatment of Alzheimer type dementia and promotion of mnemonic and learning performance.

EP 0 361 489 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 11 8014

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	CHEMICAL ABSTRACTS, vol. 88, 1978, page 621, abstract no. 22739s, Columbus, Ohio, US; Y.M. YUTILOV et al.: "New method for preparing condensed imidazolones", & KHIM. GETEROTSIKL. SOEDIN. 1976, (9), 1277-80 * N3-ethyl-3,4-pyridinediamine, RN = 61719-62-8 *	1	C 07 D 215/42 A 61 K 31/47 C 07 D 213/74 A 61 K 31/44
X	JOURNAL OF THE CHEMICAL SOCIETY, SECTION B: PHYSICAL ORGANIC CHEMISTRY, part II, 1971, pages 1425-1432, The Chemical Society, London, GB; G.B. BAR-LIN et al.: "Ionization constants of heterocyclic substances. Part IX. Protonation of aminopyridones and aminopyrimidones" * Page 1426, compounds 23,24 *	1	
A	US-A-4 734 275 (I. LALEZARI et al.)	1,8	
A	EP-A-0 258 755 (HOECHST-ROUSSEL)	1,8,9	
A	CHEMICAL ABSTRACTS, vol. 109, 12th September 1988, page 506, abstract no. 90691x, Columbus, Ohio, US C. PETERSON et al.: "Altered response of fibroblasts from aged and Alzheimer donors to drugs that elevate cytosolic free calcium", & NEUROBIOL. AGING 1988, 9(3), 261-6 * Abstract *	1,8,9	
A	CHEMICAL ABSTRACTS, vol. 110, 1989, page 15, abstract no. 205057k, Columbus, Ohio, US; P.G. WASER et al.: "4-Aminopyridine (4-AP)-derivatives as central cholinergic agents", & CURR. RES. ALZHEIMER THER.: CHOLINESTERASE INHIB. 1988, 337-42 * Abstract *	1,8,9	
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		19 March 91	DE JONG B.S.
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			